## UNITED STATES DEPARTMENT OF AGRICULTURE FOOD SAFETY AND INSPECTION SERVICE WASHINGTON. DC

## FSIS DIRECTIVE

7120.1, Revision 1

2/4/10

## SAFE AND SUITABLE INGREDIENTS USED IN THE PRODUCTION OF MEAT AND POULTRY PRODUCTS

#### I. PURPOSE

This directive provides inspection program personnel (IPP) with an up-to-date list of substances that may be used in the production of meat and poultry products. FSIS will continue to update this directive quarterly by issuing revisions to this directive as opposed to issuing amendments to the directive. This directive includes the Table of Safe and Suitable Ingredients updated as of January 5, 2010, in Amendment 21.

### II. CANCELLATION

FSIS Directive 7120.1, Safe and Suitable Ingredients Used in The Production of Meat and Poultry Products, dated August 4, 2004, and all amendments

### III. REASON FOR ISSUSSANCE

FSIS is revising this directive because it will no longer issue amendments to inspectionrelated issuances. Instead, when changes are made to FSIS directives, FSIS will reissue the entire directive as a revision. This will help ensure that inspection program personnel have all parts of directives in one document.

### IV. REFERENCES

9 CFR Chapter III

### V. BACKGROUND

A. The Table of Safe and Suitable Ingredients identifies the food grade substances that have been listed under the provisions of 9 CFR Part 424 or 21 CFR Parts 170-184 for use in meat and poultry products as food additives, generally recognized as safe substances, or other acceptable substances. This information is also available on the USDA websites at:

http://www.fsis.usda.gov/Regulations & Policies/Ingredients Guidance/index.asp

http://www.fsis.usda.gov/About FSIS/labeling & consumer protection/index.asp

**DISTRIBUTION**: Electronic **OPI**: OPPD

B. The questions and answers that follow the chart address the use of antimicrobial agents in the production of meat and poultry products.

**Assistant Administrator** 

Plity Salefler

Office of Policy and Program Development

**Table of Safe and Suitable Ingredients** 

| SUBSTANCE  | PRODUCT   | AMOUNT   | REFERENCE                   | LABELING  |
|--|---|--|-----------------------------|---|
|  |   |  |                             | REQUIREMENTS  |
|  |   | Acidifiers   |                             |   |
| Ammonium hydroxide   | pH control agent in<br>brine solutions for meat<br>products                                   | Sufficient for purpose to achieve a brine solution with a pH of 11.6 | Acceptability determination | None under the accepted conditions of use (1)                   |
| An aqueous solution of acidic calcium sulfate                              | pH control agent in<br>water used in meat and<br>poultry processing                           | Sufficient for purpose   | Acceptability determination | None under the accepted conditions of use (3)                   |
| An aqueous solution of hydrochloric and acetic acid                        | pH control agent in<br>water used in poultry<br>processing                                    | Sufficient for purpose   | Acceptability determination | None under the accepted conditions of use (3)                   |
| An aqueous solution of citric and hydrochloric acids                       | pH control agent in<br>water used in poultry<br>processing                                    | Sufficient for purpose   | Acceptability determination | None under the accepted conditions of use (1)                   |
| An aqueous solution of citric acid, hydrochloric acid, and phosphoric acid | To adjust the pH in processing water in meat and poultry plants                               | Sufficient for purpose   | Acceptability determination | None under the accepted conditions of use (1)                   |
| An aqueous solution of sulfuric acid, citric acid, and phosphoric acid     | To adjust the pH in poultry chiller water and the processing water in meat and poultry plants | Sufficient for purpose   | Acceptability determination | None under the accepted conditions of use (1)                   |
| Sodium bisulfate   | pH control agent in<br>water used in meat and<br>poultry processing                           | Sufficient for purpose   | Acceptability determination | None under the accepted conditions of use (1)                   |
| Sodium bisulfate   | pH control agent in meat and poultry soups  | Not to exceed 0.8 percent of product formulation                     | Acceptability determination | Listed by common or usual name in the ingredients statement (2) |
| Sodium bisulfate   | Added to sauces used as separable components in the formulation of various meat products      | Sufficient for purpose   | GRAS Notice<br>No. 000003   | Listed by common or usual name in the ingredients statement (2) |
| Sulfuric acid  | pH control agent in water used in poultry processing  | Sufficient for purpose   | Acceptability determination | None under the accepted conditions of use (3)                   |
|  | Ar  | nticoagulants  |                             |   |
| Sodium<br>tripolyphosphate   | Sequestrant/anti-<br>coagulant for use in<br>recovered livestock<br>blood which is            | Not to exceed 0.5 percent of recovered blood                         | Acceptability determination | Listed by common or usual name in the ingredients statement (2) |

|   | subsequently used in food products   |  |                             |   |
|---|--|--|-----------------------------|---|
|   | •  | ntimicrobials  | <u> </u>                    |   |
| An aqueous solution of sodium diacetate (4%), lactic acid (4%), pectin (2%), and acetic acid (0.5%)   | Cooked meat products   | Not to exceed 0.5 percent of finished product formulation.   | Acceptability determination | Listed by common or usual name in the ingredients statement (2) |
| An aqueous solution of sodium octanoate or octanoic acid and either glycerin and/or propylene glycol and/or a Polysorbate surface active agent (quantity sufficient to achieve the intended technical effect of octanoic acid emulsification) adjusted to a final solution pH of 1.5 to 4.0 using sodium hydroxide, potassium hydroxide, or an acceptable GRAS acid                       | Various non-<br>standardized RTE meat<br>and poultry products<br>and standardized meat<br>and poultry products<br>that permit the use of<br>any safe and suitable<br>antimicrobial agent | Applied to the surface of the product at a rate not to exceed 400 ppm octanoic acid by weight of the finished food product | Acceptability determination | None under the accepted conditions of use (3)                   |
| An aqueous solution of sodium octanoate, potassium octanoate, or octanoic acid and either glycerin and/or propylene glycol and/or a Polysorbate surface active agent (quantity sufficient to achieve the intended technical effect of octanoic acid emulsification) adjusted to a final solution pH of 1.5 to 6.0 using sodium hydroxide, potassium hydroxide, or an acceptable GRAS acid | Fresh meat primals and subprimals and cuts   | Applied to the surface of the product at a rate not to exceed 400 ppm octanoic acid by weight of the final product         | Acceptability determination | None under the accepted conditions of use (3)                   |
| A blend of citric acid<br>and sorbic acid in a 2:1<br>ratio   | To reduce the microbial load of purge trapped inside soaker pads in packages of raw whole muscle cuts of meat and poultry  | Incorporated into<br>soaker pads at a level<br>not to exceed 1 to 3<br>grams per pad                                       | Acceptability determination | None under the accepted conditions of use (1)                   |
| A blend of lactic acid (45-60%), citric acid (20-35%), and potassium hydroxide (>1%)  | Beef carcasses, beef<br>heads and beef organs  | Applied as a spray. Cannot be used on livers. All other organ meats must be drained for 1-2 minutes before                 | Acceptability determination | None under the accepted conditions of use (1)                   |

|  |  | nookoging Tolla and                        |                             | <u> </u>                              |
|--|--|--|-----------------------------|---------------------------------------|
|  |  | packaging. Tails and tongues must be       |                             |                                       |
|  |  | washed with potable                        |                             |                                       |
|  |  | water before                               |                             |                                       |
|  |  | packaging.                                 | A                           | 1.4                                   |
| A blend of salt, sodium                        | Ground beef, cooked, cured, comminuted       | Not to exceed 0.5                          | Acceptability determination | Listed by common or usual name in the |
| acetate, lemon extract, and grapefruit extract | sausages (e.g.,                              | percent of the product formulation         | determination               | ingredients                           |
| and graponale oxtract                          | bologna), and RTE                            | Torridation                                |                             | statement for the                     |
|  | whole muscle meat                            |  |                             | RTE whole muscle                      |
|  | products                                     |  |                             | meat products, and                    |
|  |  |  |                             | cooked, cured,<br>comminuted          |
|  |  |  |                             | sausages. Ground                      |
|  |  |  |                             | beef must be                          |
|  |  |  |                             | descriptively labeled                 |
|  |  |  |                             | (4)                                   |
| A blend of salt, sodium                        | Beef steaks                                  | Steaks that are sliced,                    | Acceptability determination | Product must be descriptively labeled |
| acetate, lemon extract, and grapefruit extract |  | scored and dipped in a solution containing | determination               | (4)                                   |
| and grapon an oxidati                          |  | 2.5% of the blend                          |                             | ( ' '                                 |
| A blend of salt, lemon                         | Ground beef                                  | Not to exceed 0.5                          | Acceptability               | Product must be                       |
| extract, and grapefruit                        |  | percent of the product                     | determination               | descriptively labeled                 |
| extract A blend of salt, lactic                | Various non-                                 | formulation Not to exceed 0.2              | Acceptability               | (4) All ingredients,                  |
| acid, sodium diacetate,                        | standardized RTE meat                        | percent of product                         | determination               | except for the mono-                  |
| and mono- and                                  | and poultry products                         | formulation                                |                             | and diglycerides,                     |
| diglycerides                                   | and standardized meat                        |  |                             | must be listed by                     |
|  | and poultry products that permit the use of  |  |                             | common or usual name in the           |
|  | any safe and suitable                        |  |                             | ingredients                           |
|  | antimicrobial agent                          |  |                             | statement (4)                         |
| A mixture of hops beta                         | In a salad dressing                          | Not to exceed 1.5                          | Acceptability               | Listed by common or                   |
| acids, egg white lysozyme, and cultured        | used in refrigerated meat and poultry deli   | percent of the finished salad              | determination               | usual name in the                     |
| skim milk                                      | salads                                       | Salau                                      |                             | ingredients<br>statement (2)          |
|  |  |  |                             | ( <u>-</u> )                          |
| A mixture of                                   | In salads, sauces, and                       | Not to exceed 1.5                          | Acceptability               | Listed by common or                   |
| maltodextrin (DE of 5 or                       | dressings to which fully                     | percent by weight of                       | determination               | usual name in the                     |
| greater), cultured dextrose, sodium            | cooked meat or poultry will be added         | the finished product                       |                             | ingredients<br>statement              |
| diacetate, egg white                           | Will be added                                |  |                             | (4)                                   |
| lysozyme, and nisin                            |  |  |                             | , ,                                   |
| preparation                                    | Davitaria                                    | 500 to 4000 areas 's                       | 04.050                      | Manageria                             |
| Acidified sodium chlorite                      | Poultry carcasses and parts; meat carcasses, | 500 to 1200 ppm in combination with any    | 21 CFR<br>173.325           | None under the accepted conditions    |
| Cilionite                                      | parts, meat carcasses, parts, and organs;    | GRAS acid at a level                       | 173.323                     | of use (3)                            |
|  | processed,                                   | sufficient to achieve a                    |                             | - (-/                                 |
|  | comminuted, or formed                        | pH of 2.3 to 2.9 in                        |                             |                                       |
|  | meat food products                           | accordance with 21                         |                             |                                       |
|  | (including RTE)                              | CFR 173.325 (Note:<br>The pH depends on    |                             |                                       |
|  | l  | THE PIT GOPOTIUS OIL                       | l                           | ı                                     |

|                           |   | the type of meat or poultry product.)  |  |   |
|---------------------------|---|--|--|---|
| Acidified sodium chlorite | Processed,<br>comminuted or formed<br>poultry products<br>(including RTE)                               | 500 to 1200 ppm in combination with any GRAS acid at a level sufficient to achieve a pH of 2.3 to 2.9 in accordance with 21 CFR 173.325 (Note: The pH depends on the type of meat or poultry product.)   | Acceptability determination                              | None under the accepted conditions of use (3) |
| Acidified sodium chlorite | Poultry carcasses, parts, trim, and organs  | Mixing an aqueous solution of sodium chlorite with any GRAS acid to achieve a pH of 2.2 to 3.0 then further diluting this solution with a pH elevating agent (i.e., sodium bicarbonate, sodium carbonate, or an un-acidified sodium chlorite solution) to a final pH of 5.0 to 7.5. 500 to 1200 ppm when used in a spray or dip. 50 to 150 ppm when used in a prechiller or chiller solution. Per 21 CFR 173.325.                  | Food Contact<br>Substance<br>Notification No.<br>FCN 739 | None under the accepted conditions of use (6) |
| Acidified sodium chlorite | Red meat, red meat parts and organs, and on processed, comminuted, formed meat products (including RTE) | Applied as a spray or dip, the additive is produced by mixing an aqueous solution of sodium chlorite with any GRAS acid to achieve a pH in the range of 2.2 to 3.0, then further diluting this solution with a pH elevating agent such that the resultant sodium chlorite concentration does not exceed 1200 ppm, and the chlorine dioxide concentration does not exceed 30 ppm. The pH of the use solution is between 5.0 and 7.5 | Food Contact<br>Substance<br>Notification No.<br>FCN 450 | None under the accepted conditions of use (6) |

| Ammonium hydroxide  | Beef carcasses (in hot<br>boxes and holding<br>coolers)and boneless<br>beef trimmings   | In accordance with current industry standards of good manufacturing practice  | Acceptability determination | None under the accepted conditions of use (1)   |
|---|---|---|-----------------------------|---|
| Anhydrous ammonia   | Lean finely textured<br>beef which is<br>subsequently quick<br>chilled to 28 degrees<br>Fahrenheit and<br>mechanically "stressed" | In accordance with current industry standards of good manufacturing practice  | Acceptability determination | None under the accepted conditions of use (1)   |
| Anhydrous ammonia   | Ground beef   | Followed with carbon dioxide treatment in accordance with current industry standards of good manufacturing practice                                   | Acceptability determination | None under the accepted conditions of use (1)   |
| Bacteriophage preparation (a mixture of equal proportions of six different individually purified lytic-type bacteriophages specific against <i>Listeria monocytogenes</i> ) | Various RTE meat and poultry products   | Applied as a spray at a level not to exceed 1 ml of the additive per 500 cm <sup>2</sup> product surface area   | 21 CFR<br>172.785           | Listed by common or usual name (i.e., bacteriophage preparation) in the ingredients statement of nonstandardized meat and poultry products and standardized meat and poultry products that permit the use of any safe and suitable antimicrobial agent. Standardized meat and poultry products that do not permit the use of any safe and suitable antimicrobial agent must be descriptively labeled. (4) |
| Bacteriophage preparation   | Various RTE meat and poultry products   | Applied to the surface of the product to achieve a level of 1 x 10 <sup>7</sup> to 1 x 10 <sup>9</sup> plaque forming units (pfu) per gram of product | GRAS Notice<br>No. 000218   | Listed by common or usual name (i.e., bacteriophage preparation) in the ingredients statement of nonstandardized meat and poultry products and standardized meat and poultry products that permit the use of any safe and suitable  |

|                                      | <del></del>  | T  | T                           |  |
|--------------------------------------|--|--|-----------------------------|--|
|                                      |  |  |                             | antimicrobial agent. Standardized meat and poultry products that do not permit the use of any safe and suitable antimicrobial agent must be descriptively labeled. (4) |
| Calcium hypochlorite                 | Red meat carcasses<br>down to a quarter of a<br>carcass  | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | On whole or eviscerated poultry carcasses  | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | In water used in meat processing   | Not to exceed 5 ppm calculated as free available chlorine  | Acceptability determination | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | In water used in poultry processing (except for product formulation)   | Not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | Poultry chiller water  | Not to exceed 50 ppm calculated as free available chlorine (measured in the incoming potable water)  | Acceptability determination | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | Poultry chiller red water (i.e., poultry chiller water re-circulated, usually through heat exchangers, and reused back in the chiller) | Not to exceed 5 ppm calculated as free available chlorine (measured at influent to chiller)  | Acceptability determination | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | Reprocessing contaminated poultry carcasses  | 20 ppm calculated as free available chlorine Note: Agency guidance has allowed the use of up to 50 ppm calculated as free available chlorine | 9 CFR 381.91                | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | On giblets (e.g., livers, hearts, gizzards, and necks) and salvage parts   | Not to exceed 50 ppm calculated as free available chlorine in the influent to a container for chilling.                                      | Acceptability determination | None under the accepted conditions of use (1)  |
| Calcium hypochlorite                 | Beef primals   | 20 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Carnobacterium maltaromaticum strain | Ready-to-eat comminuted meat   | Applied as a spray to meat products at a   | Gras Notice No. 000159      | All ingredients of the<br>C. maltaromaticum  |

|                          |   | 1   |                             |  |
|--------------------------|---|---|-----------------------------|--|
| CB1                      | products (e.g., hot dogs)   | maximum concentration of at inoculation of 1X10 <sup>4</sup> colony forming units per gram (cfu/g)  |                             | spray solution must<br>be listed by common<br>or usual name in the<br>ingredients<br>statement (2) |
| Cetylpyridinium chloride | To treat the surface of raw poultry carcasses prior to immersion in a chiller   | Applied as a fine mist spray of an ambient temperature aqueous solution. The aqueous solution shall also contain propylene glycol complying with 21 CFR 184.1666 at a concentration of 1.5 times that of the cetylpyridinium chloride   | 21 CFR<br>173.375           | None under the accepted conditions of use (3)  |
| Cetylpyridinium chloride | To treat the surface of raw poultry carcasses either prior to or after chilling | Not to exceed 5 gallons of solution per carcass provided that the additive is used in systems that recapture at least 99 percent of the solution that is applied to the poultry carcasses. The concentration of cetylpyridinium chloride in the solution applied to the carcasses shall not exceed 0.8 percent by weight. When application of the additive is not followed by immersion in a chiller the treatment will be followed by a potable water rinse of the carcass | 21 CFR<br>173.375           | None under the accepted conditions of use (3)  |
| Chlorine gas             | Red meat carcasses<br>down to a quarter of a<br>carcass                         | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine  | Acceptability determination | None under the accepted conditions of use (1)  |
| Chlorine gas             | On whole or eviscerated poultry carcasses                                       | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine  | Acceptability determination | None under the accepted conditions of use (1)  |
| Chlorine gas             | In water used in meat processing  | Not to exceed 5 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Chlorine gas             | In water used in poultry  | Not to exceed 50 ppm  | Acceptability               | None under the   |

|                  | product formulation)   | available chlorine  |                             | of use (1)  |
|------------------|--|---|-----------------------------|---|
| Chlorine gas     | Poultry chiller water  | Not to exceed 50 ppm calculated as free available chlorine (measured in the incoming potable water)   | Acceptability determination | None under the accepted conditions of use (1)                   |
| Chlorine gas     | Poultry chiller red water (i.e., poultry chiller water re-circulated, usually through heat exchangers, and reused back in the chiller) | Not to exceed 5 ppm calculated as free available chlorine (measured at influent to chiller)   | Acceptability determination | None under the accepted conditions of use (1)                   |
| Chlorine gas     | Reprocessing contaminated poultry carcasses  | 20 ppm calculated as free available chlorine Note: Agency guidance has allowed the use of up to 50 ppm calculated as free available chlorine  | 9 CFR 381.91                | None under the accepted conditions of use (1)                   |
| Chlorine gas     | On giblets (e.g., livers, hearts, gizzards, and necks) and salvage parts   | Not to exceed 50 ppm calculated as free available chlorine in the influent to a container for chilling.   | Acceptability determination | None under the accepted conditions of use (1)                   |
| Chlorine gas     | Beef primals   | 20 ppm calculated as free available chlorine  | Acceptability determination | None under the accepted conditions of use (1)                   |
| Citric acid      | Bologna in an edible casing  | Up to a 10 percent solution applied prior to slicing  | Acceptability determination | Listed by common or usual name in the ingredients statement (4) |
| Citric acid      | Bologna in an inedible casing  | Up to a 10 percent solution applied prior to slicing  | Acceptability determination | None under the accepted conditions of use (1)                   |
| Citric acid      | Fully cooked meat and poultry products in impermeable casings.   | Up to a 3 percent solution is applied to the casing just prior to removal.  | Acceptability determination | None under the accepted conditions of use (1)                   |
| Citric acid      | Separated beef heads<br>and associated offal<br>products (e.g., hearts,<br>livers, tails, tongues)                                     | A 2.5 percent solution applied as a spray prior to chilling   | Acceptability determination | None under the accepted conditions of use (1)                   |
| Chlorine dioxide | In water used in poultry processing  | Not to exceed 3 ppm residual chlorine dioxide as determined by Method 4500-ClO <sub>2</sub> E in the "Standard Methods for the Examination of Water and Wastewater," 18 <sup>th</sup> | 21 CFR<br>173.300           | None under the accepted conditions of use (3)                   |

|  |  | ed., 1992, or an equivalent method  |   |  |
|--|--|---|---|--|
| Chlorine dioxide                                   | Red meat, red meat<br>parts and organs;<br>processed,<br>comminuted, or formed<br>meat food products   | Applied as a spray or dip at a level not to exceed 3 ppm residual chlorine dioxide as determined by Method 4500-ClO <sub>2</sub> E in the "Standard Methods for the Examination of Water and Wastewater," 18 <sup>th</sup> ed., 1992, or an equivalent method | Food Contact<br>Substance<br>Notification No.<br>FCN 668                | None under the accepted conditions of use (6)  |
| Cultured Sugar (derived from corn, cane, or beets) | In enhanced meat and poultry products (e.g., beef or pork injected with a solution) and RTE meat and poultry products (e.g., hot dogs and cooked turkey breast)                            | At up to 4.8 percent of the product formula   | GRAS Notice<br>No. 000240   | Cultured cane and beet sugar listed by common or usual name (e.g., "cultured cane sugar) or as "cultured sugar."  Cultured corn sugar listed as "cultured corn sugar" or "cultured dextrose" (2) |
| DBDMH (1,3dibromo-<br>5,5-dimethylhydantoin)       | For use in poultry chiller water and in water applied to poultry via an Inside-Outside Bird Washer (IOBW) and in water used in poultry processing for poultry carcasses, parts, and organs | At a level not to exceed that needed to provide the equivalent of 100 ppm active bromine  | Food Contact<br>Substance<br>Notification No.<br>FCN 334 and<br>FCN 453 | None under the accepted conditions of use (6)  |
| DBDMH (1,3dibromo-<br>5,5-dimethylhydantoin)       | For use in water supplied to ice machines to make ice intended for general use in poultry processing   | At a level not to exceed that needed to provide the equivalent of 100 ppm of available bromine (corresponding to a maximum level of 90 mg DBDMH/kg water)   | Food Contact<br>Substance<br>Notification No.<br>FCN 775                | None under the accepted conditions of use (6)  |
| DBDMH (1,3dibromo-<br>5,5-dimethylhydantoin)       | For use in water applied to beef hides, carcasses, heads, trim, parts, and organs.   | At a level not to exceed that needed to provide the equivalent of 300 ppm active bromine.   | Food Contact<br>Substance<br>Notification No.<br>FCN 792                | None under the accepted conditions of use (6)  |
| Egg white lysozyme                                 | In casings and on cooked (RTE) meat and poultry products   | 2.5 mg per pound in<br>the finished product<br>when used in casings;<br>2.0 mg per pound on   | GRAS Notice<br>No. 000064   | Listed by common or usual name in the ingredients statement (2)  |

|   |  | cooked meat and  |                             |   |
|---|--|--|-----------------------------|---|
| Electrolytically<br>generated<br>hypochlorous acid                                  | Red meat carcasses<br>down to a quarter of a<br>carcass  | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1) |
| Electrolytically generated hypochlorous acid  | On whole or eviscerated poultry carcasses  | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1) |
| Electrolytically<br>generated<br>hypochlorous acid                                  | In water used in meat processing   | Not to exceed 5 ppm calculated as free available chlorine  | Acceptability determination | None under the accepted conditions of use (1) |
| Electrolytically<br>generated<br>hypochlorous acid                                  | In water used in poultry processing (except for product formulation)   | Not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1) |
| Electrolytically generated hypochlorous acid  | Poultry chiller water  | Not to exceed 50 ppm calculated as free available chlorine (measured in the incoming potable water)  | Acceptability determination | None under the accepted conditions of use (1) |
| Electrolytically<br>generated<br>hypochlorous acid                                  | Poultry chiller red water (i.e., poultry chiller water re-circulated, usually through heat exchangers, and reused back in the chiller) | Not to exceed 5 ppm calculated as free available chlorine (measured at influent to chiller)  | Acceptability determination | None under the accepted conditions of use (1) |
| Electrolytically<br>generated<br>hypochlorous acid                                  | Reprocessing contaminated poultry carcasses  | 20 ppm calculated as free available chlorine Note: Agency guidance has allowed the use of up to 50 ppm calculated as free available chlorine | 9 CFR 381.91                | None under the accepted conditions of use (1) |
| Electrolytically<br>generated<br>hypochlorous acid                                  | On giblets (e.g., livers, hearts, gizzards, and necks) and salvage parts   | Not to exceed 50 ppm calculated as free available chlorine in the influent to a container for chilling.                                      | Acceptability determination | None under the accepted conditions of use (1) |
| Electrolytically<br>generated<br>hypochlorous acid                                  | Beef primals   | 20 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1) |
| An aqueous solution of citric and hydrochloric acids adjusted to a pH of 1.0 to 2.0 | Poultry carcasses, parts, trim, and organs   | Applied as a spray or dip with a minimum contact time of 2 to 5 seconds  | Acceptability determination | None under the accepted conditions of use (1) |
| An aqueous solution of citric and hydrochloric acids adjusted to a pH of 0.5 to 2.0 | Meat carcasses, parts, trim, and organs  | Applied as a spray or dip for a contact time of 2 to 5 seconds   | Acceptability determination | None under the accepted conditions of use (1) |
| A blend of citric acid  | Poultry carcasses  | Applied as a spray   | Acceptability               | None under the                                |

| (1.87%), phosphoric<br>acid (1.72%), and<br>hydrochloric acid<br>(0.8%)  |  | with a minimum contact time of 1 to 2 seconds and allowed to drip from the carcasses for 30 seconds                                | determination               | accepted conditions of use (1)  |
|--|--|--|-----------------------------|---|
| A blend of citric acid,<br>hydrochloric acid, and<br>phosphoric acid   | To adjust the acidity in various meat and poultry products   | Sufficient for purpose   | Acceptability determination | Listed by common or usual name in the ingredients statement (2)   |
| Hops beta acids  | In casings and on cooked (RTE) meat and poultry products   | 2.5 mg per pound in<br>the finished product<br>when used in casings;<br>2.0 mg per pound on<br>cooked meat and<br>poultry products | GRAS Notice<br>No. 000063   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)  |
| Lactic Acid  | Livestock carcasses<br>prior to fabrication (i.e.,<br>pre- and post-chill),<br>offal, and variety meats                    | Up to a 5 percent lactic acid solution   | Acceptability determination | None under the accepted conditions of use (1)   |
| Lactic acid  | Beef and pork sub-<br>primals and trimmings  | 2 percent to 5 percent<br>solution of lactic acid<br>not to exceed 55°C  | Acceptability determination | None under the accepted conditions of use (1)   |
| Lactic acid  | Beef heads and tongues   | A 2.0 to 2.8 percent solution applied to brushes in a washer cabinet system used to clean beef heads and tongues                   | Acceptability determination | None under the accepted conditions of use (1)   |
| Lactic acid bacteria<br>mixture consisting of<br>Lactobacillus<br>acidophilus (NP35,<br>NP51), Lactobacillus<br>lactis (NP7), and<br>Pediococcus acidilactici<br>(NP3) | RTE cooked sausages<br>(e.g., frankfurters,<br>bologna, etc.) and<br>cooked, cured whole<br>muscle products (e.g.,<br>ham) | Applied by dipping product into a solution containing 10 <sup>7</sup> colony forming units lactobacilli per ml                     | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)  |
| Lactic acid bacteria<br>mixture consisting of<br>Lactobacillus<br>acidophilus (NP35,<br>NP51), Lactobacillus<br>lactis (NP7), and<br>Pediococcus acidilactici<br>(NP3) | Poultry carcasses and fresh whole muscle cuts and chopped/ground poultry   | 10 <sup>5</sup> to 10 <sup>6</sup> colony<br>forming units of<br>lactobacilli per gram of<br>product                               | Acceptability determination | Listed by common or usual name in the ingredients statement of nonstandardized products. Single ingredient raw products must be descriptively labeled (2) |
| Lactic acid bacteria<br>mixture consisting of<br>Lactobacillus<br>acidophilus (NP35,<br>NP51), Lactobacillus   | Non-standardized<br>comminuted meat<br>products (e.g., beef<br>patties), ground beef,<br>and raw whole muscle              | 10 <sup>6</sup> to 10 <sup>8</sup> colony<br>forming units of<br>lactobacilli per gram of<br>product                               | GRAS Notice<br>No. 000171   | Listed by common or<br>usual name in the<br>ingredients<br>statement of non-<br>standardized  |

| lactis (NP7), and Pediococcus acidilactici (NP3)  Lactoferrin               | Beef carcasses and parts   | At up to 2 percent of a water-based antimicrobial spray  | GRAS Notice<br>No. 000067   | comminuted meat products. Ground beef and raw whole muscle beef cuts must be descriptively labeled (2) Listed by common or usual name in ingredients statement (2)                               |
|---|--|--|-----------------------------|--|
| Lactoferrin   | Beef carcasses   | As part of an antimicrobial spray that would deliver 1 gram of lactoferrin per dressed beef carcass, followed by a wash with tempered water and rinse with lactic acid | GRAS Notice<br>No. 000130   | None under the accepted conditions of use (1)  |
| Lauramide arginine ethyl ester (LAE), silicon dioxide, and refined sea salt | Non-standardized RTE comminuted meat products and standardized RTE comminuted meat products that permit the use of any safe and suitable antimicrobial agent   | Not to exceed 200<br>ppm LAE by weight of<br>the finished product  | Acceptability determination | Listed by common or usual name (i.e., lauric arginate, refined sea salt) in the ingredients statement (2)  |
| Lauramide arginine ethyl ester (LAE),                                       | Fresh cuts of meat and poultry; and, non-  | Not to exceed 200<br>ppm LAE, 67 ppm   | Acceptability determination | Listed by common or usual name (i.e.,  |
| silicon dioxide, and refined sea salt                                       | standardized, non-<br>comminuted RTE<br>meat and poultry<br>products and<br>standardized, non-<br>comminuted RTE<br>meat and poultry<br>products that permit<br>the use of any safe<br>and suitable<br>antimicrobial agent | silicon dioxide, and<br>1640 ppm refined<br>sea salt by weight of<br>the finished product  |                             | lauric arginate, silicon dioxide, refined sea salt) in the ingredients statement (2) When applied to the surface of fresh cuts of meat and poultry none under the accepted conditions of use (1) |

| Eauramide arginine ethyl ester (LAE) dissolved at specified concentrations in either propeling and various flower products are retained to the concentrations in either propeline glycol, glycerin, or water to a standardized RTE meat and poultry products that permit the use of active agent (quantity sufficient to achieve the intended technical effect of LAE emulsification)    Lauramide arginine ethyl ester (LAE)   RTE meat and poultry products; raw pork sausage   Applied to the inside of the package via a process known as "Sprayed Lethality in Container" (SLIC) or similar process at up to 44 ppm (with a process tolerance of 20 percent, allowing for an LAE concentration not to exceed 53 ppm) by weight of the finished product state permit the use of an acceptability and the surface of RTE meat and poultry products.    Nisin preparation   Cooked, RTE meat and poultry soups   Not to exceed 50 pm of the product formulation   Nisin preparation   In casings and on cooked (RTE) meat and poultry products.    Nisin preparation   Nisin preparation   Nisin preparation   Cooked (RTE) meat and poultry products   Statement (2)   Not to exceed 50 pm of the product formulation   Cooked (RTE) meat and poultry products   Statement (2)   CRAS Notice   Not to exceed 50 pm of the product formulation   CRAS Notice   Not to exceed 50 pm of the product formulation   CRAS Notice   Not to exceed 50 pm of the product formulation   CRAS Notice   Not to exceed 50 pm of the product formulation   CRAS Notice   Not to exceed 50 pm of the product formulation   CRAS Notice   Not to exceed 50 pm of the product formulation   CRAS Notice   CRAS Notice   Not to exceed 50 pm of the product formulation   CRAS Notice   CRAS Notice   Not to exceed 50 pm of the product   CRAS Notice   Not to exceed 50 pm of the product   CRAS Notice   Not to exceed 50 pm of the product   CRAS Notice   Not to exceed 50 pm of the product   CRAS Notice   CRAS Notice   Not to exceed 50 pm of the product   CRAS Notice   Not to exceed 50 pm of the product   CRAS Notice   Not t   | emulsification)   |   |  |                             |  |
|--|---|---|--|-----------------------------|--|
| the package via a process known as "Sprayed Lethality in Container" (SLIC) or similar process tolerance of 20 percent, allowing for an LAE concentration not to exceed 53 ppm) by weight of the finished product containing sauces  Nisin preparation  Nisin preparation  Nisin preparation  Not to exceed 5 ppm of the product formulation  Not to exceed 55 ppm of the product formulation  Not to exceed 55 ppm of the product formulation  Not to exceed 55 ppm of the product formulation  A blend of encapsulated nisin preparation (90.9) percent), rosemary extract (8.2 percent) and salt (0.9 percent)  A blend of finsin  Cooked (RTE) meat and poultry sausages  Prankfurters and other similar cooked meat and poultry sausages  Not to exceed 550 ppm of the product when used in casings; 2.5 mg per pound on cooked meat and poultry products  A blend of finsin  Cooked (RTE) meat and other similar cooked meat and poultry sausages  Not to exceed 550 ppm of the product when used in casings; 2.5 mg per pound on cooked meat and poultry products  Not to exceed 550 ppm of the product formulation  A coeptability determination or usual name in the ingredients statement (2)  Listed by common or usual name in the ingredients statement (2)  Listed by common or usual name in the ingredients statement (2)  A coeptability determination  A coep | Lauramide arginine ethyl ester (LAE) dissolved at specified concentrations in either propylene glycol, glycerin, or water to which may be added a Polysorbate surface active agent (quantity sufficient to achieve the intended technical effect of LAE | poultry and various<br>non-standardized RTE<br>meat and poultry<br>products and<br>standardized RTE meat<br>and poultry products<br>that permit the use of<br>any safe and suitable | of the product at a rate<br>not to exceed 200<br>ppm LAE by weight of<br>the finished food   |                             | surface of RTE products listed by common or usual name (i.e., lauric arginate) in the ingredients statement (2) When applied to the surface of fresh cuts of meat and poultry none under the accepted conditions |
| Nisin preparation  |   | products; raw pork  | the package via a process known as "Sprayed Lethality in Container" (SLIC) or similar process at up to 44 ppm (with a process tolerance of 20 percent, allowing for an LAE concentration not to exceed 53 ppm) by weight of the finished |                             | None under the accepted conditions   |
| Nisin preparation  In casings and on cooked (RTE) meat and poultry products  A blend of encapsulated nisin preparation (90.9 percent), rosemary extract (8.2 percent) and salt (0.9 percent)  A blend of nisin  Soups  of the product formulation  of the product formulation  of the product formulation  of the product formulation  determination  GRAS Notice No. 000065  No. 000065  Listed by common or usual name in the ingredients statement (2)  Not to exceed 550 ppm of the product formulation  A cceptability determination  usual name in the ingredients statement (2)  Listed by common or usual name in the ingredients statement (2)  Not to exceed 550 ppm of the product formulation  A blend of nisin  Cooked (RTE) meat  Not to exceed 0.55  Acceptability  Listed by common or usual name in the ingredients statement (4)   | Nisin preparation   | poultry products  | ppm nisin preparation  |                             | ingredients  |
| Cooked (RTE) meat and poultry products  A blend of encapsulated nisin preparation (90.9 percent), rosemary extract (8.2 percent) and salt (0.9 percent)  A blend of nisin  Cooked (RTE) meat the finished product when used in casings; 2.5 mg per pound on cooked meat and poultry products  No. 000065  Acceptability determination  Listed by common or usual name in the ingredients statement (2)  A blend of nisin  Cooked (RTE) meat  No. 000065  Acceptability  Acceptability  Listed by common or usual name in the ingredients statement (4)  Acceptability  Listed by common or usual name in the ingredients statement (4)  Acceptability  Listed by common or usual name in the ingredients statement (4)   | Nisin preparation   |   | of the product   |                             | or usual name in the ingredients   |
| encapsulated nisin preparation (90.9 percent), rosemary extract (8.2 percent) and salt (0.9 percent)  A blend of nisin  similar cooked meat and poultry sausages  ppm of the product formulation  ppm of the product formulation  determination usual name in the ingredients statement (4)  statement (4)  Not to exceed 0.55  Acceptability  Listed by common or   |   | cooked (RTE) meat   | the finished product<br>when used in<br>casings; 2.5 mg per<br>pound on cooked<br>meat and poultry<br>products   |                             | Listed by common<br>or usual name in<br>the ingredients  |
|  | encapsulated nisin<br>preparation (90.9<br>percent), rosemary<br>extract (8.2 percent)<br>and salt (0.9 percent)  | similar cooked meat<br>and poultry sausages   | ppm of the product formulation   | determination               | usual name in the ingredients statement (4)  |
| ANTANARANDO ROCEMBRO I ANA DAUTRO CALICARDE LI DERCANT OT NOMBRO LI MATARMINATION LI LICUAL NAMA IN TOA  | A blend of nisin preparation, rosemary  | and poultry sausages  | Not to exceed 0.55 percent of product  | Acceptability determination | Listed by common or usual name in the  |

| An aqueous mixture of peroxyacetic acid, hydrogen peroxide, acetic acid, and 1-   | carcasses as a spray, wash, rinse, dip, chiller water, or scald water  (1) Water or ice for washing, rinsing, cooling, or otherwise processing whole or                           | In either application,<br>the level of<br>peroxyacetic acid<br>will not exceed 220  | Food Contact<br>Substance<br>Notification<br>No. FCN           | None under the accepted conditions of use (6)                            |
|---|---|---|--|--|
| A mixture of peroxyacetic acid, hydrogen peroxide, acetic acid, and 1-hydroxyethylidene-1, 1-diphosphonic acid (HEDP)     | (1) Process water for washing, rinsing, cooling, or otherwise for processing meat carcasses, parts, trim, and organs; and (2) process water applied to poultry parts, organs, and | In either application,<br>the level of<br>peroxyacetic acid<br>will not exceed 230<br>ppm, hydrogen<br>peroxide will not<br>exceed 165 ppm, and<br>HEDP will not exceed<br>14.0 ppm   | Food Contact<br>Substance<br>Notification<br>No. FCN<br>000323 | None under the accepted conditions of use (6)                            |
| octanoic acid, acetic acid, hydrogen peroxide, peroxyoctanoic acid, and 1-hydroxyethylidene-1, 1-diphosphonic acid (HEDP) | carcasses, parts, trim and organs   | concentrations for meat carcasses, parts, and organs: Peroxyacetic acids 220 ppm, hydrogen peroxide 75 ppm; Maximum concentrations for poultry carcasses, parts, and organs: Peroxyacetic acids 220 ppm, hydrogen peroxide 110 ppm, HEDP 13 ppm | 173.370  | accepted conditions of use (3)   |
| Ozone  Peroxyacetic acid,   | All meat and poultry products  Meat and poultry   | In accordance with current industry standards of good manufacturing practice  Maximum   | 21 CFR<br>173.368  | None under the accepted conditions of use (3)  None under the            |
| Organic Acids (i.e., lactic, acetic, and citric acid)   | As part of a carcass wash applied pre-chill   | At up to 2.5 percent of a solution  | FSIS Notice 49-<br>94  | None under the accepted conditions of use (1)                            |
| A blend of nisin<br>preparation, rosemary<br>extract, salt, and<br>sodium diacetate                                       | Cooked (RTE) meat<br>and poultry sausages<br>and cured meat<br>products   | ppm)  Not to exceed 0.25 percent of product formulation (where the nisin preparation will not exceed 250 ppm)   | Acceptability determination                                    | Listed by common or<br>usual name in the<br>ingredients<br>statement (4) |
| extract, salt,<br>maltodextrin, and<br>cultured dextrose  | and cured meat products   | formulation in cooked (RTE) meat and poultry sausages and 0.7 percent of product formulation in cured meat products (where the nisin preparation will not exceed 250  |  | ingredients<br>statement (4)   |

| 1-diphosphonic acid  | parts, trim, and   | peroxide will not   |   |  |
|--|--|---|---|--|
| (HEDP) and optionally sulfuric acid  | organs; and, (2) water or ice applied to whole or cut poultry including parts, trim, and organs as a spray, wash, rinse, dip, chiller water or scalder water                           | exceed 85 ppm, and<br>HEDP will not exceed<br>11 ppm.   |   |  |
| A mixture of Peroxyacetic acid and 1-hydroxyethylidene-1, 1-diphosphonic acid (HEDP)   | Poultry finishing chillers<br>and in process water<br>applied to poultry parts,<br>organs, and carcasses<br>in low temperature<br>(e.g., less than 40<br>degrees F) immersion<br>baths | In either application,<br>the level of<br>Peroxyacetic acid will<br>not exceed 2000 ppm<br>and HEDP will not<br>exceed 136 ppm  | Food Contact<br>Substance<br>Notification No.<br>FCN 000880 | None under the accepted conditions of use (6)  |
| Potassium diacetate  | Various meat and poultry products which permit the addition of antimicrobial agents, e.g., hot dogs  | Not to exceed 0.25 percent of the product formulation   | Acceptability determination                                 | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)   |
| A solution of water, lactic acid, propionic acid, and acidic calcium sulfate (solution with a pH range of 1.0-2.0)*                    | Various RTE meat products, e.g., hot dogs.   | Applied as a spray for 20-30 seconds of continual application just prior to packaging *Propionic acid may be removed from the solution; sodium phosphate may be added to the solution as a buffering agent (the amount of sodium phosphate on the finished product must not exceed 5000ppm. | Acceptability determination                                 | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)   |
| A solution of water,<br>acidic calcium sulfate<br>and 85-95,000 ppm of<br>lactic acid (solution with<br>a pH range of 0.35 to<br>0.55) | Raw comminuted beef.   | To treat raw beef during grinding to lower the pH of the product.   | Acceptability determination                                 | Product must be descriptively labeled (2)  |
| A solution of water, acidic calcium sulfate, lactic acid, and sodium phosphate (solution with a pH range of 1.45 to 1.55)              | Raw whole muscle beef cuts and cooked roast beef and similar cooked beef products (e.g., corned beef, pastrami, etc.).   | Spray applied for up to 30 seconds of continual application *sodium phosphate on the finished product must not exceed 5000 ppm.   | Acceptability determination                                 | Listed by common or usual name in the ingredients statement of multi-ingredient products. Single ingredient roast beef products and raw whole muscle beef cuts must be descriptively labeled (2) |

| A solution of water, acidic calcium sulfate, lactic acid, and sodium phosphate (solution with a pH of 1.45 to 1.6)  | Cooked poultry carcasses and parts.   | Spray applied for 20 to 40 seconds of continual application * sodium phosphate on the finished product must not exceed 5000 ppm. | Acceptability determination | Listed by common or usual name in the ingredients statement of multi-ingredient products. Single ingredient whole muscle cuts of poultry must be descriptively labeled |
|---|---|--|-----------------------------|--|
| A solution of water, acidic calcium sulfate, lactic acid, and disodium phosphate (solution with a pH of 1.0 to 2.0) | Beef jerky  | Applied to the surface of the product with a contact time not to exceed 30 seconds   | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)   |
| Skim milk or dextrose cultured with propionibacterium freudenreichii subsp. Shermanii                               | Meat and poultry sausages including those with standards of identity which permit the use of antimicrobial agents | Not to exceed 2 percent by weight of the finished product  | GRAS Notice<br>No. 000128   | Listed by common or usual name in the ingredients statement (2)  |
| Sodium citrate buffered with citric acid to a pH of 5.6   | Non-standardized and standardized comminuted meat and poultry products which permit ingredients of this type      | Not to exceed 1.3 percent of the product formulation in accordance with 21 CFR 184.1751  | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)   |
| Sodium hypochlorite   | Red meat carcasses<br>down to a quarter of a<br>carcass   | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Sodium hypochlorite   | On whole or eviscerated poultry carcasses   | Applied as a spray at a level not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Sodium hypochlorite   | In water used in meat processing  | Not to exceed 5 ppm calculated as free available chlorine  | Acceptability determination | None under the accepted conditions of use (1)  |
| Sodium hypochlorite   | In water used in poultry processing (except for product formulation)  | Not to exceed 50 ppm calculated as free available chlorine   | Acceptability determination | None under the accepted conditions of use (1)  |
| Sodium hypochlorite   | Poultry chiller water   | Not to exceed 50 ppm calculated as free available chlorine (measured in the incoming potable water)                              | Acceptability determination | None under the accepted conditions of use (1)  |
| Sodium hypochlorite   | Poultry chiller red water (i.e., poultry chiller water re-circulated, usually through heat exchangers, and        | Not to exceed 5 ppm calculated as free available chlorine (measured at influent to chiller)                                      | Acceptability determination | None under the accepted conditions of use (1)  |

|                                | reused back in the chiller)  |   |  |   |
|--------------------------------|--|---|--|---|
| Sodium hypochlorite            | Reprocessing contaminated poultry carcasses                              | 20 ppm calculated as free available chlorine Note: Agency guidance has allowed the use of up to 50 ppm calculated as free available chlorine  | 9 CFR 381.91   | None under the accepted conditions of use (1)                   |
| Sodium hypochlorite            | On giblets (e.g., livers, hearts, gizzards, and necks) and salvage parts | Not to exceed 50 ppm calculated as free available chlorine in the influent to a container for chilling.   | Acceptability determination                                | None under the accepted conditions of use (1)                   |
| Sodium hypochlorite            | Beef primals   | 20 ppm calculated as free available chlorine  | Acceptability determination                                | None under the accepted conditions of use (1)                   |
| Sodium metasilicate            | Component of marinades used for raw meat and poultry products            | Not to exceed 2 percent by weight of the marinade   | Acceptability determination                                | None under the accepted conditions of use (1)                   |
| Sodium metasilicate            | Raw beef carcasses, subprimals, and trimmings                            | A 4 percent (plus or minus 2 percent) solution  | Acceptability determination                                | None under the accepted conditions of use (1)                   |
| Sodium metasilicate            | RTE meat products  | Up to a 6 percent solution applied to the surface of the product at a rate not to exceed 300 ppm of the finished product  | Acceptability determination                                | None under the accepted condition of use (1)                    |
| Trisodium phosphate            | Raw unchilled poultry carcasses and giblets                              | 8-12 percent solution applied by spraying or dipping giblets for up to 30 seconds. 8-12 percent solution within a temperature range of 65° F to 85° F applied by spraying or dipping carcasses for up to 15 seconds | Acceptability<br>determination<br>(per 21 CFR<br>182.1778) | None under the accepted conditions of use (1)                   |
|                                | F  | ntioxidants   |  |   |
| BHA (butylated hydroxyanisole) | "Brown N Serve"<br>sausages  | 0.02 percent in combination with other antioxidants for use in meat, based on fat content   | Acceptability determination                                | Listed by common or usual name in the ingredients statement (4) |
| BHT (butylated hydroxytoluene) | "Brown N Serve"<br>sausages  | 0.02 percent in combination with other antioxidants for use in meat, based on fat content   | Acceptability determination                                | Listed by common or usual name in the ingredients statement (4) |

|   |   | Binders   |                             |   |
|---|---|---|-----------------------------|---|
| A mixture of sodium<br>alginate, calcium<br>sulfate, glucono delta-<br>lactone, and sodium<br>pyrophosphate | Various meat and poultry products where binders are permitted   | Mixture not to exceed 1.55 percent of product formulation with the sodium alginate not to exceed 1 percent of the product formulation and the sodium pyrophosphate not to exceed 0.5 percent of the product formulation | Acceptability determination | Listed by common or usual name in the ingredients statement (4)             |
| A mixture of carrageenan, whey protein concentrate, and xanthan gum   | Sausages where binders are permitted; cooked poultry products; beef and poultry patties; modified breakfast sausage, cooked sausages, and fermented sausages covered by FSIS Policy Memo 123; and modified substitute versions of fresh sausage, ground beef, or hamburger covered by FSIS Policy Memo 121B | Not to exceed 3.5 percent by weight of the product formulation  | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement<br>(4) |
| Beef collagen   | Various meat and poultry products where binders are permitted   | Not to exceed 3.5 percent of product formulation  | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement<br>(4) |
| Binders listed in 9 CFR<br>424.21(c) for use in<br>cured pork products<br>and poultry products              | "Turkey ham and water products"   | In accordance with 9<br>CFR 319.104(d) and<br>424.21(c)   | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)    |
| Carboxymethyl cellulose (cellulose gum)   | Poultry franks  | Not to exceed 3.5 percent of the product formulation  | Acceptability determination | Listed by common or usual name in the ingredients statement (4)             |
| Carboxymethyl cellulose   | Cured pork products   | Not to exceed 3 percent of product formulation  | Acceptability determination | Listed by common or usual name in the ingredients statement (4)             |
| Carrot Fiber  | Various comminuted meat and poultry products where binders are permitted  | Not to exceed 3.5 percent of the product formulation  | GRAS Notice<br>No. 000116   | List as "isolated<br>carrot product"<br>(2)                                 |
| Cellulose, powdered conforming to the specifications in the   | Various comminuted poultry products where binders are permitted   | Not to exceed 3.5 percent of the product formulation  | Acceptability determination | Listed by common or usual name in the ingredients                           |

| Food Chemicals Codex 5 <sup>th</sup> Edition |  |  |  | statement (2)   |
|--|--|--|--|---|
| Guar powder,<br>micronized                   | Various meat and poultry products where binders are permitted  | Not to exceed 3.0 percent of the product formulation   | Acceptability determination  | Listed by common or usual name in the ingredients statement (4) |
| Hydroxypropyl<br>methylcellulose             | Seasoning mixtures added to sauces and gravies produced under FDA jurisdiction that will be used in meat and poultry products  | Sufficient for purpose   | Acceptability determination  | Listed by common or usual name in the ingredients statement (2) |
| Hydroxypropyl<br>methylcellulose             | Thickener in meat and poultry pot pie fillings, sauces, soups, and gravies   | Not to exceed 1 percent of the product formulation   | Acceptability determination  | Listed by common or usual name in the ingredients statement (2) |
| Inulin                                       | Various meat and poultry products (e.g., frankfurters, sausage, patties, loaves, pates) where binders are permitted  | 2 to 5 percent of the product formulation  | Acceptability<br>determination<br>and GRAS<br>Notice No.<br>000118 | Listed by common or usual name in the ingredients statement (2) |
| Konjac flour                                 | Meat and poultry products in which starchy vegetable flours are permitted  | No to exceed 3.5 percent of the product formulation individually or collectively with other binders  | Acceptability determination  | Listed by common or usual name in the ingredients statement (2) |
| Methylcellulose                              | Various comminuted meat and poultry products where binders are permitted   | Not to exceed 3.5 percent of the product formulation   | Acceptability determination  | Listed by common or usual name in the ingredients statement (2) |
| Methylcellulose                              | Thickener in meat and poultry pot pie fillings, sauces, soups, and gravies; a binder in poultry patties, loaves, and nuggets; a binder in meat patties, loaves, and nuggets; texturizer in Policy Memo 121B and 123 products | Not to exceed 1 percent of the product formulation as a thickener in meat and poultry pot pie fillings, sauces, soups, and gravies; 1.6 percent as a binder in poultry patties, loaves, and nuggets; 0.25 percent as a binder in meat patties, loaves, and nuggets; 0.6 percent as a texturizer in Policy Memo 121B and 123 products | Acceptability determination  | Listed by common or usual name in the ingredients statement (2) |
| Oat Hull Fiber                               | Various non-<br>standardized<br>comminuted meat<br>products  | Not to exceed 3.5 percent of the product formulation   | GRAS Notice<br>No. 000261  | Listed as "isolated oat product" in the ingredients statement   |
| Oat Fiber                                    | Various meat products  | Not to exceed 3.5  | Acceptability  | Listed as "isolated   |

|   | (e.g., frankfurters, sausage patties, loaves) where binders are permitted and whole muscle meat products  | percent of the product formulation                   | determination               | oat product" in the ingredients statement. Whole muscle meat products must be descriptively labeled (4) |
|---|---|--|-----------------------------|---|
| Orange pulp, dried                                      | Non-standardized whole muscle meat and poultry products where binders are permitted and standardized whole muscle meat and poultry products where standards of identity permit the use of binders | Not to exceed 3.5 percent of the product formulation | Acceptability determination | List as "citrus<br>flour" or "dried<br>orange pulp" (2)   |
| Orange pulp, dried and orange pulp, dried with guar gum | Various ground meat<br>and poultry products<br>where binders are<br>permitted   | Not to exceed 3.5 percent of the product formulation | GRAS Notice<br>No. 000154   | List as "citrus flour"<br>or "dried orange<br>pulp" (2)   |
| Partially hydrolyzed proteins                           | Various meat and poultry products where binders are permitted.  | Not to exceed 3.5 percent of the product formulation | Acceptability determination | Listed by common or usual name in the ingredients statement (2)   |
| Pectin  | Various meat and poultry products where binders are permitted   | Not to exceed 3 percent of the product formulation   | Acceptability determination | Listed by common or usual name in the ingredients statement (2)   |
| Pork collagen   | Various meat and poultry food products where binders are permitted  | Not to exceed 3.5 percent of the product formulation | Acceptability determination | Listed by common or usual name in the ingredients statement (2)   |
| Pork skin proteins                                      | Various meat products where binders are permitted   | Not to exceed 1.5 percent of product formulation     | Acceptability determination | Listed by common or usual name in the ingredients statement (4)   |
| Rice bran   | Various comminuted<br>meat and poultry<br>products where binders<br>are permitted (e.g., hot<br>dogs, meatballs, and<br>chicken patties)  | Not to exceed 3.5 percent of the product formulation | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)                                |
| Rice starch   | Cured pork products   | Not to exceed 0.8 percent of product formulation     | Acceptability determination | Listed by common or usual name in the ingredients statement (4)   |
| Sodium alginate   | Various meat products where binders are permitted   | Not to exceed 1 percent of the product formulation   | Acceptability determination | Listed by common or usual name in the ingredients statement (2)   |
| Sodium alginate   | Various poultry   | Not to exceed 0.8                                    | Acceptability               | Listed by common or   |

|  | products where binders are permitted  | percent of the product formulation  | determination               | usual name in the ingredients statement (2)                              |
|--|---|---|-----------------------------|--|
| "(species) protein" (e.g., chicken protein)                              | Whole muscle poultry food products where binders are permitted provided the protein is used in products of the same kind (e.g., chicken protein in a marinade injected into whole muscle chicken food products) | Not to exceed 0.225 percent of the marinade solution  | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (2) |
| "(species) protein" (e.g., chicken protein, concentrated turkey protein) | Various poultry products where the protein solution is used in products of the same kind (e.g., chicken protein in a coating of a breaded chicken fritter)  | As a coating applied to the product and/or as a portion of the batter. Not to exceed 0.8 percent of product formulation when applied as a protein coating only, 0.14 percent of product formulation when used in the batter only, and 0.89 percent of product formulation when used as both a coating and in the batter | GRAS Notice<br>No. 000168   | Listed by common or usual name in the ingredients statement (2)          |
| Transglutaminase enzyme  | Texturizing agent in meat and poultry food products where texturizing agents and binders are permitted  | Not to exceed 65 ppm<br>of the product<br>formulation   | Acceptability determination | Listed by common or usual name in the ingredients statement (2)          |
| Transglutaminase enzyme  | Cross-linking agent in modified meat and poultry products addressed in Policy Memos 121B and 123.   | Not to exceed 65 ppm<br>of the product<br>formulation   | Acceptability determination | Listed by common or usual name in the ingredients statement (2)          |
| Transglutaminase enzyme  | Binding and cross-<br>linking agent in<br>uncooked restructured<br>chicken breasts  | Not to exceed 100 ppm of the product formulation  | Acceptability determination | Listed by common or usual name in the ingredients statement (2)          |
| Trehalose  | Binding and purge<br>control agent in various<br>meat and poultry<br>products where binders<br>are permitted  | Not to exceed 2<br>percent of the product<br>formulation  | GRAS Notice<br>No. 000045   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2) |
| Xanthan gum (purified by recovery with ethyl                             | Various meat and poultry products where   | Non-standardized meat and poultry   | GRAS Notice<br>No. 000121   | Listed by common or usual name in the                                    |

| alaahal)                          | hindoro oro normitto d                | products and products                        |                             | ingradianta                    |
|-----------------------------------|---------------------------------------|--|-----------------------------|--------------------------------|
| alcohol)                          | binders are permitted                 | products and products with a standard of     |                             | ingredients<br>statement (4)   |
|                                   |                                       | identity which currently                     |                             | Statement (4)                  |
|                                   |                                       | permit the use of                            |                             |                                |
|                                   |                                       | xanthan gum listed in                        |                             |                                |
|                                   |                                       | 9 CFR 424.21(c)                              |                             |                                |
|                                   | Cc                                    | oloring Agents                               |                             |                                |
| Carmine (cochineal)               | To color isolated soy                 | 0.2 to 0.4 percent of                        | Acceptability               | Listed by common or            |
|                                   | protein for use in dry                | the hydrated protein                         | determination               | usual name in the              |
|                                   | cured acidified                       | gel. The protein gel                         |                             | ingredients                    |
|                                   | sausages                              | must not exceed 30                           |                             | statement (5)                  |
|                                   |                                       | percent of the meat                          |                             |                                |
|                                   |                                       | food product                                 |                             |                                |
| Ourina Assala                     |                                       | formulation                                  |                             |                                |
|                                   | erators (must be us                   |  |                             |                                |
| Potassium erythorbate             | Cured pork and beef                   | 87.5 oz. to 100 gallons                      | Acceptability               | Listed by common or            |
|                                   | cuts; cured meat food products; cured | of pickle at 10 percent pump; 7/8 oz. to 100 | determination               | usual name in the ingredients  |
|                                   | comminuted poultry or                 | lbs. Of meat, meat                           |                             | statement (2)                  |
|                                   | poultry products                      | byproduct or poultry                         |                             | Statement (2)                  |
|                                   | poditry products                      | product; 10 percent to                       |                             |                                |
|                                   |                                       | surfaces of cured                            |                             |                                |
|                                   |                                       | meat cuts or poultry                         |                             |                                |
|                                   |                                       | products prior to                            |                             |                                |
|                                   |                                       | packaging                                    |                             |                                |
|                                   |                                       |  |                             |                                |
| Denuding agent<br>by rinsing with | ts (may be used in potable water.)    | combination. Mu                              | ist be remov                | ed from tripe                  |
| Calcium carbonate                 | Denuding agent for                    | Sufficient for purpose                       | Acceptability               | None under the                 |
|                                   | washing tripe                         |  | determination               | accepted conditions            |
|                                   |                                       |  |                             | of use (1)                     |
| Calcium citrate                   | Denuding agent for                    | Sufficient for purpose                       | Acceptability               | None under the                 |
|                                   | washing tripe                         |  | determination               | accepted conditions            |
| Calairma brednassida              | Danishing agent for                   | Cufficient for numero                        | A a a a m t a b i lite :    | of use (1)                     |
| Calcium hydroxide                 | Denuding agent for washing tripe      | Sufficient for purpose                       | Acceptability determination | None under the                 |
|                                   | wasning tripe                         |  | determination               | accepted conditions of use (1) |
| Potassium carbonate               | Denuding agent for                    | Sufficient for purpose                       | Acceptability               | None under the                 |
| 1 otassiam carbonate              | washing tripe                         | Sumsient for purpose                         | determination               | accepted conditions            |
|                                   | washing tripe                         |  | determination               | of use (1)                     |
| Potassium citrate                 | Denuding agent for                    | Sufficient for purpose                       | Acceptability               | None under the                 |
|                                   | washing tripe                         |  | determination               | accepted conditions            |
|                                   |                                       |  |                             | of use (1)                     |
| Potassium hydroxide               | Denuding agent for                    | Sufficient for purpose                       | Acceptability               | None under the                 |
|                                   | washing tripe                         |  | determination               | accepted conditions            |
| T2: 12: 12: 12: 12: 1             | D                                     | 0.000  | A ( - 1 - 12)               | of use (1)                     |
| Tricalcium phosphate              | Denuding agent for                    | Sufficient for purpose                       | Acceptability               | None under the                 |
|                                   | washing tripe                         |  | determination               | accepted conditions            |
| Tripotassium                      | Denuding agent for                    | Sufficient for purpose                       | Acceptability               | of use (1)  None under the     |
| phosphate                         | washing tripe                         | ounicient for purpose                        | determination               | accepted conditions            |
| priospriate                       | washing inpe                          |  | determination               | accepted conditions            |

|  |  |   |                             | of use (1)  |
|--|--|---|-----------------------------|---|
|  | Film   | Forming Agents  |                             |   |
| A mixture of water,<br>glycerin, carrageenan,<br>and cornstarch  | Used to aid in the release of elastic netting on cooked meat products that are cooked in elastic netting | Sufficient for purpose  | Acceptability determination | None under the accepted conditions of use (1)                                       |
| A mixture of water,<br>glycerin, carageenan,<br>cornstarch, and<br>caramel   | Used to aid in the release of elastic netting on cooked meat products that are cooked in elastic netting | Sufficient for purpose  | Acceptability determination | "Caramel Color" listed as an ingredient and as a product name qualifier (2)         |
| A mixture of water,<br>glycerin, carageenan,<br>cornstarch, and smoke<br>flavoring   | Used to aid in the release of elastic netting on cooked meat products that are cooked in elastic netting | Sufficient for purpose  | Acceptability determination | "Smoke Flavor" listed<br>as an ingredient and<br>as a product name<br>qualifier (2) |
| A solution of sodium alginate, dextrose, isolated pea protein, sugar, and maltodextrin (DE of 6) used with a solution of calcium chloride, powdered sugar, oleoresin black pepper, and isolated pea protein. | Used to form a calcium alginate-based casing on pork and poultry sausages.                               | Quantity of the casing on the sausage ranges from 8 to 15 percent of total product formulation and calcium alginate not to exceed 0.219 percent of the finished product formulation | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (4)            |
| Gelatin spice sheets   | To ensure even distribution of seasonings on cooked pork products  | Sufficient for purpose  | Acceptability determination | None under the accepted conditions of use (1)                                       |
| Hydroxypropyl<br>methylcellulose   | Film-forming agent in glazes for meat and poultry products   | Not to exceed 4 percent of the product formulation  | Acceptability determination | Listed by common or usual name in the ingredients statement (2)                     |
| Methylcellulose  | Film-forming agent in glazes for meat and poultry products   | Not to exceed 3 percent of the product formulation for poultry products, 3.5 percent of the product formulation for meat products   | Acceptability determination | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)            |
|  | Fla  | voring Agents   |                             |   |
| Adenosine 5'- monophosphoric acid (AMP) and its monosodium and disodium salts  | As a flavor enhancer for meat and poultry soups and soup mixes   | Not to exceed 200 ppm of the product formulation  | GRAS Notice<br>No. 000144   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)            |

| Lactic acid   | As a flavor enhancer added to pork fatty tissue used in the production of dehydrated pork fatty tissue                       | Not to exceed 0.367 percent of the pork fatty tissue, prior to dehydration   | Acceptability determination                                    | Product must be descriptively labeled (4)   |
|---|--|--|--|---|
| Laminaria japonica<br>(brown algae)                                   | As a flavor enhancer or flavoring agent in marinades for meat and poultry, meat and poultry soups, gravies, and seasonings   | Not to exceed 0.08 percent of the product formulation  | GRAS Notice<br>No. 000123                                      | Listed by common or usual name in the ingredients statement (2)                                   |
| Potassium acetate   | Various meat and poultry products  | No to exceed 1.2 percent of the product formulation  | Acceptability determination                                    | Listed by common or usual name in the ingredients statement (4)                                   |
| Sucralose   | Non-nutritive sweetener in various non-standardized meat and poultry products  | Not to exceed 500 ppm in the product formulation   | Acceptability determination                                    | Listed by common or usual name in the ingredients statement (2)                                   |
| Trehalose   | As a flavor enhancer in non-standardized RTE meat and poultry products   | Not to exceed 2 percent by weight of product formulation   | Acceptability determination                                    | Listed by common or usual name in the ingredients statement (2)                                   |
|   | M  | iscellaneous   |  |   |
| Alkyl polyglycosides  | Hog scalding   | Sufficient for purpose of increasing the wetting ability of the caustic solution   | GRAS Notice<br>No. 000237                                      | None under the accepted conditions of use (1)   |
| Alkyl polyglycosides  | Wash meat (i.e., beef carcasses after the hide has been removed to remove any extraneous hair, dirt, etc.) during butchering | Used at 2% active solution level followed by a potable water rinse   | GRAS Notice<br>No. 000237                                      | None under the accepted conditions of use (1)   |
| Ammonium hydroxide  | To adjust the pH of brine solutions prior to injection into meat   | Sufficient for purpose to achieve a brine solution with a pH of up to 11.6   | Acceptability determination                                    | None under the accepted conditions of use (1)   |
| An aqueous solution of arginine, potassium hydroxide, salt, and water | pH control agent in<br>brine solutions for<br>beef subprimals or to<br>make beef patties                                     | Arginine is added to the salt and water brine solution at 0.2-0.6% by weight and the pH is maintained between 10.4 and 10.7. The potassium hydroxide is then added at 0.0528 – 0.0793% by weight to adjust the pH to 11.4 – 11.9 | Acceptability determination L-arginine: GRAS Notice No. 000290 | Salt and water must<br>be listed by<br>common or usual<br>name on the<br>ingredients<br>statement |
| A 60/40 blend of sodium bicarbonate                                   | To generate carbon dioxide in packages of  | Incorporated into soaker pads at a level   | Acceptability determination                                    | None under the accepted conditions  |

| and citric acid  | raw whole muscle cuts  | not to exceed 0.5 to 2   |                             | of use (1)  |
|--|--|--|-----------------------------|---|
| A solution of water, dextrose, glycerin, maltose, and sodium phosphate | of meat and poultry  To aid in the removal of residual blood from beef and bison carcasses after the typical exsanguination process is completed | grams per pad Sufficient for purpose   | Acceptability determination | For all edible tissue none under the accepted conditions of use unless the Moisture Fat Free% (MFF%) analysis shows treated carcasses are not in compliance with retained water requirements. All edible tissue from treated carcasses not in compliance must be labeled in accordance with Policy Memo 066C. Organ meat from all treated carcasses must be descriptively labeled (1) |
| Algal oil derived from Schizochytrium sp.                              | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 1.45 percent by weight of the product formulation for meat products and 0.87 percent by weight of the product formulation for poultry products | GRAS Notice<br>No. 000137   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)  |
| Cellulose (powdered)   | To facilitate grinding and shredding in cheese   | No to exceed 2 percent of the cheese   | Acceptability determination | None under the accepted conditions of use (1)   |
| Citroglycerides (citric acid esters of monoand diglycerides)           | To aid in the dispersion of lauric arginate (LAE)  | Used in a 5:1 mixture with lauric arginate with the maximum amount in meat and poultry products not to exceed 1125 ppm                                       | GRAS Notice<br>No. 000222   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)  |
| Cultured Sugar (derived from cane, corn, or beets)                     | In uncooked (raw) sausage meat   | At up to 4.8 percent of the product formula  | GRAS Notice<br>No. 000240   | Cultured cane and beet sugar listed by common or usual name (e.g., "cultured cane sugar) or as "cultured sugar."  Cultured corn sugar listed as "cultured corn sugar" or "cultured dextrose" (2)  |

| Diacylglycerol oil  | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 11 percent of the meat or poultry product formula  | GRAS Notice<br>No. 000115   | Listed by common or usual name in the ingredients statement (2)          |
|---|--|--|-----------------------------|--|
| Erythorbic Acid   | To delay discoloration in ground beef and ground beef patties                                | Not to exceed 0.04 percent of the product formulation  | Acceptability determination | Product must be descriptively labeled (2)                                |
| Fish oil concentrate  | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 2.9 percent by weight of the product formulation for meat products and 1.7 percent by weight of the product formulation for poultry products | GRAS Notice<br>No. 000105   | Listed by common or usual name in the ingredients statement (2)          |
| Fish oil (predominantly sardine, anchovy, and tuna)   | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 3.3 percent by weight of the product formulation for meat products and 2.0 percent by weight of the product formulation for poultry products | GRAS Notice<br>No. 000193   | Listed by common or usual name in the ingredients statement (2)          |
| Fish oil (predominantly anchovy)  | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 3.3 percent by weight of the product formulation for meat products and 2.0 percent by weight of the product formulation for poultry products | GRAS Notice<br>No. 000138   | Listed by common or usual name in the ingredients statement (2)          |
| Fish oil (predominantly anchovy) microencapsulated  | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 6.0 percent by weight of the product formulation for meat products and 3.6 percent by weight of the product formulation for poultry products | GRAS Notice<br>No. 000138   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2) |
| Glucose oxidase and catalase enzymes from Aspergillus niger with a dextrose energy source and sodium bicarbonate buffer | To maintain a low oxygen atmosphere in packages of raw whole muscle cuts of meat and poultry | Incorporated into<br>soaker pads such that<br>the enzymes do not<br>exceed 0.03 percent<br>by weight of the meat<br>or poultry                             | Acceptability determination | None under the accepted conditions of use (1)                            |
| Glucose oxidase and catalase enzymes from Aspergillus niger with a  | To maintain a low oxygen atmosphere in packages of shelf-                                    | Applied to the surface of the product such that the enzymes do   | Acceptability determination | Listed by common or usual name in the ingredients                        |

| dextrose energy source<br>and sodium<br>bicarbonate buffer  | stable, ready-to-eat,<br>meat products   | not exceed 0.03 percent by weight of the meat food product   |                             | statement (2)   |
|---|--|--|-----------------------------|---|
| Glycerophospholipid cholesterol acyltransferase (GCAT) enzyme preparation from Bacillus licheniformis expressing a modified GCAT gene from Aeromonas salmonicida subsp. salmonicida (GCAT enzyme preparation) | For use as an emulsifier in comminuted meat products   | Not to exceed 22.6 mg TOS/kg of total product formulation  | GRAS Notice<br>No. 000265   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)  |
| Hydrogen peroxide   | To minimize biofilm buildup on reverse osmosis and ultrafiltration membranes for processing beef plasma  | Not to exceed 100 ppm added just prior to plasma entering membranes  | Acceptability determination | None under the accepted conditions of use (1)   |
| Hydrolyzed gelatin  | To prevent moisture loss from fresh cuts of meat and poultry   | A 13 percent aqueous solution of hydrolyzed gelatin sprayed on the surface not to exceed 2 percent hydrolyzed gelatin by weight of the meat or poultry | Acceptability determination | Listed by common or usual name in the ingredients statement. Label must also bear a statement, contiguous to the product name, indicating product has been coated with hydrolyzed gelatin to prevent moisture loss. (4) |
| Medium and long chain triacylglycerol (tailored triglycerides containing approximately 12 percent medium chain fatty acids)   | For use as a supplementary source of vegetable oil in the production of various meat and poultry products  | Sufficient for purposes  | GRAS Notice<br>No. 000217   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)  |
| Polyglycerol ester produced by transesterification of triglycerol with soybean oil  | Added to fresh livestock blood during collection to eliminate foaming  | Not to exceed 8.8 ppm<br>in the fresh livestock<br>blood   | Acceptability determination | None under the accepted conditions of use (1)   |
| Polyglycerol<br>polyricinoleic acid<br>(PGPR)   | For use as an emulsifier in the formulation of color additives which are subsequently used in processed meat and poultry products for which colors are | Sufficient for purpose using good manufacturing practices  | GRAS Notice<br>No. 000270   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)  |

|  | permitted  |  |                             |  |
|--|--|--|-----------------------------|--|
| Salmon oil   | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 5.0 percent by weight of the product formulation for meat products and 3.0 percent by weight of the product formulation for poultry products | GRAS Notice<br>No. 000146   | Listed by common or<br>usual name in the<br>ingredients<br>statement (2) |
| Small planktivorous pelagic fish oil                       | For use as an alternative edible oil in the production of various meat and poultry products  | Not to exceed 3.3 percent by weight of the product formulation for meat products and 2.0 percent by weight of the product formulation for poultry products | GRAS Notice<br>No. 000102   | Listed by common or usual name in the ingredients statement (2)          |
| Sodium bicarbonate   | Neutralize excess<br>acidity (maintain pH) in<br>fresh pork and beef<br>cuts   | In an injected solution,<br>not to exceed 0.5<br>percent of the product<br>formulation   | Acceptability determination | Listed by common or usual name in the ingredients statement (2)          |
| Sodium bicarbonate   | Maintain pH and reduce purge in fresh turkey products  | In an injected solution,<br>not to exceed 0.5<br>percent of the product<br>formulation   | Acceptability determination | Listed by common or usual name in the ingredients statement (2)          |
| Sodium bicarbonate   | To soak natural casings to ease stuffing   | 1.06 percent of an aqueous solution. Casings must be rinsed with potable water prior to stuffing   | Acceptability determination | None under the accepted conditions of use (1)                            |
| Sodium hydroxide   | For application to poultry carcasses immediately after removal of feathers and prior to evisceration to minimize fecal material from adhering to the carcass | 0.05 percent solution  | Acceptability determination | None under the conditions of use (1)                                     |
| Sodium hydroxide and hydrochloric acid                     | To adjust the pH of (species) plasma during processing (in which it is exposed to heat) to prevent gelling   | Sufficient for purpose to adjust pH  | Acceptability determination | None under the accepted conditions of use (1)                            |
| Stearidonic acid<br>(SDA) soybean oil                      | For use as an ingredient in meat and poultry products  | Sufficient for purpose   | GRAS Notice<br>No. 000283   | Listed by common or usual name in the ingredients statement (2)          |
| Triple salt of magnesium, ammonium, and potassium chloride | For use as a substitute for a portion of the sodium chloride normally used in meat   | Sufficient for purpose   | GRAS Notice<br>No. 000272   | Listed by common or usual name in the ingredients statement (2)          |

|   | and poultry products.   |   |                                |  |
|---|---|---|--------------------------------|--|
| Trisodium phosphate (as a component of phosphate blends, not to exceed 40 percent of the phosphate blend)                             | To decrease the amount of cooked out juices in meat food products except where otherwise prohibited by the meat inspection regulations and poultry food products except where otherwise prohibited by the poultry products inspection regulations | For meat food products, 5 percent of phosphate in pickle at 10 percent pump level; 0.5 percent of phosphate in meat food product (only clear solution may be injected into meat food product). For poultry food products, 0.5 percent of total product.                               | Acceptability determination    | Listed by common or usual name in the ingredients statement (4) Note: Phosphates may be collectively designated as "sodium phosphates" or "potassium phosphates" |
| Tuna oil  | For use as an alternative edible oil in the production of various meat and poultry products   | Not to exceed 3.1 percent by weight of the product formulation for meat products and 1.8 percent by weight of the product formulation for poultry products  | GRAS Notice<br>No. 000109      | Listed by common or<br>usual name in the<br>ingredients<br>statement (2)   |
| Xanthan gum   | Suspending agent for carrageenan in a brine tank  | Not to exceed 2 percent of the amount of carrageenan  | Acceptability determination    | None under the accepted conditions of use (1)  |
|   | Pack  | aging Systems   |                                |  |
| Carbon monoxide gas<br>as part of Cryovac's<br>modified atmosphere<br>packaging system<br>(for use with 550P<br>Tray/Lid and LID551P) | Packaging fresh cuts of case ready muscle meat and case ready ground meat to maintain wholesomeness, provide flexibility in distribution, and reduce shrinkage of the meat  | The use of carbon monoxide (0.4 percent), carbon dioxide (30 percent) and nitrogen (69.6 percent) as part of the Cryovac low oxygen modified atmosphere packaging system used with 550P Tray /Lid   | Acceptability<br>Determination | None under the accepted conditions of use (2)  |
| Carbon monoxide gas<br>as part of Cryovac's<br>modified atmosphere<br>packaging system  | Packaging fresh cuts of case ready muscle meat and case ready ground meat to maintain wholesomeness   | The use of carbon monoxide (0.4 percent), carbon dioxide (30 percent) and nitrogen (69.6 percent) introduced directly into the package. System uses a barrier lid that only covers a highly permeable patch. The permeable patch is a one half inch hole in the lid film. Barrier lid | Acceptability determination    | None under the accepted conditions of use (2)  |

|   |   | removed prior to display for retail sale  |                             |   |
|---|---|---|-----------------------------|---|
| Carbon monoxide gas<br>as part of the Pactiv<br>modified atmosphere<br>packaging system<br>(ActiveTech 2001)  | Packaging fresh cuts of case ready muscle meat and case ready ground meat to maintain wholesomeness                       | The use of carbon monoxide (0.4 percent), carbon dioxide (30 percent) and nitrogen (69.6 percent) as part of the Pactiv modified atmosphere packaging system  | GRAS Notice<br>No. 000083   | None under the accepted conditions of use (2)   |
| Carbon monoxide gas<br>as part of a high<br>oxygen modified<br>atmosphere packaging<br>system used in<br>accordance with GRN<br>000083 (Cargill)  | Packaging fresh cuts of case-ready muscle meat and ground meat to maintain wholesomeness                                  | Not to exceed 0.4 percent of the modified atmosphere gas mixture  | Acceptability determination | None under the accepted conditions of use (2)   |
| Carbon monoxide gas a part of Cargill's modified atmosphere packaging system introduced directly into the bulk or master container used for bulk transportation of fresh meat products. Meat products are subsequently repackaged in packages not containing a carbon monoxide modified atmosphere prior to retail sale (In accordance with GRN 000083) | Packaging fresh cuts of muscle meat and ground meat to maintain wholesomeness   | Not to exceed 0.4 percent of the modified atmosphere gas mixture  | Acceptability determination | None under the accepted conditions of use (2)   |
| Carbon monoxide gas<br>as part of the Precept<br>modified atmosphere<br>packaging system  | Packaging case-ready<br>fresh cuts of beef and<br>pork as well as ground<br>beef and pork to<br>maintain<br>wholesomeness | Carbon monoxide 0.4 percent (with a process tolerance of 20 percent, allowing for a carbon monoxide concentration up to 0.48 percent) in combination with carbon dioxide (20-100 percent) and nitrogen (0-80 percent) | GRAS Notice<br>No. 000143   | None under the accepted conditions of use (2) Products packaged in this MAP system must be coded with a "Use or Freeze by" date not to exceed 28 days after packaging for ground meat and 35 days for whole muscle cuts |
| Carbon monoxide gas<br>as part of Precept's<br>modified atmosphere<br>packaging system  | Packaging case-ready<br>fresh cuts of poultry as<br>well as ground poultry  | Carbon monoxide 0.3 percent (with a process tolerance of 20 percent, allowing for a carbon monoxide   | Acceptability determination | None under the accepted conditions of use (2) Products packaged in this MAP system  |

| Carbon monoxide as a component of a modified atmosphere packaging system (Tyson Foods, Inc.) | Packaging case-ready<br>fresh cuts of beef and<br>pork as well as ground<br>beef and pork | concentration up to 0.36 percent), in combination with nitrogen (0-80 percent), and carbon dioxide (20-100 percent)  Carbon monoxide (at a level not to exceed 2.2 mg carbon monoxide per pound of packaged meat) in combination with carbon dioxide and nitrogen | GRAS Notice<br>No. 000167   | must be coded with a "Use or Freeze by" date not to exceed 28 days after packaging for ground poultry and 35 days for whole muscle cuts of poultry  None under the accepted conditions of use (2) Products packaged in this MAP system must be coded with a "Use or Freeze by" date not to exceed 28 days after packaging for ground meat and 35 days for |
|--|---|---|-----------------------------|---|
| Poultry scald a  | gents (must be re   | moved by subse  | uont cloanir                | whole muscle cuts   |
| Alkyl polyglycosides   | To remove feathers  | Sufficient for purpose  | GRAS Notice                 | None under the  |
| Aikyi polyglycosides   | from poultry carcasses  | Summer for purpose  | No. 000237                  | conditions of use (1)   |
| Calcium acid   | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| phosphate  | from poultry carcasses  | Cumoletti for purpose   | determination               | conditions of use (1)   |
| Calcium acid   | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| pyrophosphate  | from poultry carcasses  |   | determination               | conditions of use (1)   |
| Calcium bicarbonate  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
|  | from poultry carcasses  | ' '   | determination               | conditions of use (1)   |
| Calcium carbonate  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
|  | from poultry carcasses  |   | determination               | conditions of use (1)   |
| Calcium  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| dodecylbenzene   | from poultry carcasses  |   | determination               | conditions of use (1)   |
| sulfonate  |   |   |                             |   |
| Calcium 2-ethylhexyl   | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| sulfate  | from poultry carcasses  |   | determination               | conditions of use (1)   |
| Calcium  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| hexametaphosphate  | from poultry carcasses  |   | determination               | conditions of use (1)   |
| Calcium hydroxide  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
|  | from poultry carcasses  | 0 (" : (  | determination               | conditions of use (1)   |
| Calcium lauryl sulfate   | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| Coloium phoophata  | from poultry carcasses  | Cufficient for purpose  | determination               | conditions of use (1)   |
| Calcium phosphate (mono-, di-, and   | To remove feathers  | Sufficient for purpose  | Acceptability determination | None under the conditions of use (1)  |
| tribasic)  | from poultry carcasses  |   | ueterriniation              | Conditions of use (1)   |
| Calcium pyrophosphate  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| - Saloidin pyrophosphale   | from poultry carcasses  | Cumolonic for purpose   | determination               | conditions of use (1)   |
| Calcium  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
| sesquicarbonate  | from poultry carcasses  | 2 3   | determination               | conditions of use (1)   |
| Calcium sulfate  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |
|  | from poultry carcasses  |   | determination               | conditions of use (1)   |
| Calcium  | To remove feathers  | Sufficient for purpose  | Acceptability               | None under the  |

| tripolyphosphate          | from poultry carcasses |                        | determination    | conditions of use (1) |
|---------------------------|------------------------|------------------------|------------------|-----------------------|
| Potassium acid            | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| phosphate                 | from poultry carcasses |                        | determination    | conditions of use (1) |
| Potassium acid            | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| pyrophosphate             | from poultry carcasses |                        | determination    | conditions of use (1) |
| Potassium bicarbonate     | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
|                           | from poultry carcasses |                        | determination    | conditions of use (1) |
| Potassium carbonate       | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
|                           | from poultry carcasses |                        | determination    | conditions of use (1) |
| Potassium                 | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| dodecylbenzene            | from poultry carcasses |                        | determination    | conditions of use (1) |
| sulfonate                 |                        |                        |                  | , ,                   |
| Potassium 2-ethylhexyl    | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| sulfate                   | from poultry carcasses |                        | determination    | conditions of use (1) |
| Potassium                 | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| hexametaphosphate         | from poultry carcasses |                        | determination    | conditions of use (1) |
| Potassium hydroxide       | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| ,                         | from poultry carcasses | Cameron Fare           | determination    | conditions of use (1) |
| Potassium lauryl sulfate  | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| . otacolam ladry rodinate | from poultry carcasses | Camelelli lei pai pece | determination    | conditions of use (1) |
| Potassium phosphate       | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| (mono-, di-, and          | from poultry carcasses | Camelelli lei pai pece | determination    | conditions of use (1) |
| tribasic)                 | mom pounty cardadeed   |                        | dotorriiridation |                       |
| Potassium                 | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| pyrophosphate             | from poultry carcasses | Camelon to parpood     | determination    | conditions of use (1) |
| Potassium                 | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| sesquicarbonate           | from poultry carcasses | Camelon to parpood     | determination    | conditions of use (1) |
| Potassium sulfate         | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| 1 otassiam sanate         | from poultry carcasses | Cambient for purpose   | determination    | conditions of use (1) |
| Potassium                 | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| tripolyphosphate          | from poultry carcasses | Camelon to parpood     | determination    | conditions of use (1) |
| Tetracalcium              | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| pyrophosphate             | from poultry carcasses | Camelon to parpood     | determination    | conditions of use (1) |
| Tetrapotassium            | To remove feathers     | Sufficient for purpose | Acceptability    | None under the        |
| pyrophosphate             | from poultry carcasses | Cambient for purpose   | determination    | conditions of use (1) |
| ругорпоорпасо             |                        | dorizina Agonto        | dotorriiridation |                       |
|                           |                        | derizing Agents        | T                | Ι                     |
| Calcium gluconate         | Raw meat products      | Solutions applied or   | Acceptability    | Listed by common or   |
|                           |                        | injected into raw meat | determination    | usual name in the     |
|                           |                        | shall not result in a  |                  | ingredients           |
|                           |                        | gain of 3 percent      |                  | statement (2)         |
|                           |                        | above green weight     |                  |                       |
| Protease preparation      | Raw meat products      | Solutions applied or   | Acceptability    | Listed by common or   |
| derived from Bacillus     |                        | injected into raw meat | determination    | usual name in the     |
| subtilis                  |                        | shall not result in a  |                  | ingredients           |
|                           |                        | gain of 3 percent      |                  | statement (2)         |
|                           |                        | above green weight     |                  |                       |
| Protease produced         | Raw meat products      | Solutions applied or   | Acceptability    | Listed by common or   |
| from Bacillus subtilis    |                        | injected into raw meat | determination    | usual name in the     |
| var. amyloliquefaciens    |                        | shall not result in a  |                  | ingredients           |
|                           |                        | gain of 3 percent      |                  | statement (2)         |
|                           |                        | above green weight     |                  |                       |
| Protease produced         | Raw meat cuts and raw  | Solutions applied or   | GRAS Notice      | Listed by common or   |

| from Aspergillus niger | poultry muscle tissue of | injected into raw meat    | No. 000089 | usual name in the |
|------------------------|--------------------------|---------------------------|------------|-------------------|
|                        | hen, cock, mature        | or poultry tissue shall   |            | ingredients       |
|                        | turkey, mature duck,     | not result in a gain of 3 |            | statement (2)     |
|                        | mature goose, and        | percent above green       |            |                   |
|                        | mature guinea            | weight                    |            |                   |

- 1) The use of the substance(s) is consistent with FDA's labeling definition of a processing aid.
- 2) Generally Recognized as Safe (GRAS)
- 3) Secondary Direct Food Additive
- 4) Direct Food Additive
- 5) Color Additive
- 6) Food Contact Substance
- \* Substances identified in bold print in the table are substances that have been added to the directive since it was last issued on December 17, 2002.

## Questions and Answers on the Use of Antimicrobial Agents in the Production of Meat and Poultry Products

The following set of questions and answers provide information regarding the requirements for the use of antimicrobial agents in meat and poultry production.

#### References

-Final Rule, "Food Ingredients and Sources of Radiation Listed or Approved for Use in the Production of Meat and Poultry Products" (December 1999).

- -MOU between FDA and FSIS for Ingredient Approval (January, 2000).
- -FSIS Directive 7120.1, "Safe and Suitable Ingredients Used in the Production of Meat and Poultry Products."
- -Guidance document on "Ingredients and Sources of Radiation Used to Reduce Microorganisms on Carcass, Ground Beef and Beef Trimmings."
- --Guidance Procedures for Notification and Protocol Submission of New Technology, February 2004 <a href="http://www.fsis.usda.gov/regulations">http://www.fsis.usda.gov/regulations</a> & policies/New Technology Notification & Protocol Submissi on/index.asp
- -Federal Register Notice, "FSIS Procedures for Notification of New Technology" (68 FR 6873) (February, 2003)
- -9 CFR Part 416.4
- -FSIS Directive 6355.1, "Use of Chlorine Dioxide in Poultry Chill Water."
- -9 CFR 424.21(c)
- FSIS Directive 6700.1 and <u>FSIS Directive 6700.1 Amendment 1</u> <u>Html</u>, "Retained Water in Raw Meat and Poultry Products."
- -21 CFR Part 172,173, 182, 184
- -21 CFR 101.100 (a)(3)(ii)(c)

### 1. Question: What is the definition of a New Technology?

Answer: According to the FSIS Federal Register Notice (68 FR 6873) entitled, "FSIS Procedures for Notification of New Technology," FSIS defines a "new technology" as new, or new applications of, equipment, substances, methods, processes or procedures affecting the slaughter of livestock and poultry or processing of meat, poultry, or egg products which could affect product safety, inspection procedures, inspection program personnel safety, or require a waiver of a regulation.

#### 2. Question: What is the definition of a processing aid?

Answer: According to the Food and Drug Administration's (FDA) regulations (21 CFR 101.100 (a) (3) (ii)), the definition of a processing aid is:

- a. Substances that are added to a food during the processing of such food but are removed in some manner from the food before it is packaged in its finished form.
- b. Substances that are added to a food during processing, are converted into constituents normally present in the food, and do not significantly increase the amount of the constituents naturally found in food.
- c. Substances that are added to a food for their technical or functional effect in the processing but are present in the finished food at insignificant levels and do not have any technical or functional effect in that food.

An example of a processing aid is the use of organic acid(s) (e.g., lactic, acetic, or citric acid) as part of a livestock carcass wash applied pre-chill.

FSIS has posted guidelines on processing aids in regulating the labeling of meat and poultry products at:

http://www.fsis.usda.gov/PDF/Determination\_of\_Processing\_Aids.pdf.

http://www.fsis.usda.gov/PDF/Prohibited\_Substances\_in\_FSIS\_Actions\_on%20\_Use\_of\_Ingredient\_s.pdf

#### 3. Question: What are secondary direct food additives and direct food additives?

Answer: According to FDA's regulations (21 CFR Part 173), secondary direct food additives are substances whose functionality is required during the manufacture or processing of a food and are ordinarily removed from the final food. Although residuals might carry over to the final food, residuals must not exhibit any technical effects. Secondary direct food additives are consistent with FDA's definition of a processing aid so labeling is not required. Examples of secondary direct food additives are acidified sodium chlorite (21 CFR 173.325) and peroxyacids (21 CFR 173.370).

According to FDA's regulations (21 CFR Part 172), direct food additives are used to provide a technical effect in the final food. The antioxidants BHA and BHT are examples of substances that are approved as direct food additives.

## 4. Question: Do organic acid(s) (e.g., lactic, acetic, or citric acid) that are used as antimicrobial agents need to be declared on the label if they are applied to livestock carcasses after the chilling step?

Answer: Organic acid(s) are generally recognized as safe (GRAS) and are listed in FSIS regulations for use as an acidifier in various meat and poultry products at a level which is sufficient for purpose (9 CFR 424.21(c)). All ingredients, including organic acid(s), require labeling unless the use of a substance is consistent with FDA's definition of a processing aid or is a secondary direct food additive.

FSIS has recently stated no objection to the use of 5% hot lactic acid as an antimicrobial agent to treat beef carcasses prior to fabrication (i.e., pre and post-chill). Data was submitted to the Agency that demonstrated no lasting effect under the specified conditions of use. FSIS determined that the proposed use is consistent with the definition of a processing aid. Therefore, its use would not need to be reflected on the labeling for treated carcasses or products produced from treated carcasses. This new use is listed in the table of this directive.

If a company is interested in using one or more of these organic acid(s) as an antimicrobial agent on livestock carcasses or trim in a manner other than which is currently approved, they must provide data to the Agency that show that the use complies with FDA's definition of a processing aid. The data must show that the organic acid has only a momentary technical effect, not a lasting effect on the meat, e.g., fresh color is not preserved, normal spoilage indicators (e.g. discoloration) are not masked; and there is no extension of shelf life as compared to products made with untreated trimmings. The data must also show that the nutrient composition is not affected by the treatment and the sensory characteristics of the product are not affected. (Note: the reference to "Guidance on Ingredients and Sources of Radiation used to Reduce Microorganisms on Carcasses, Ground Beef, and Beef Trim," can be accessed at http: <a href="https://www.fsis.usda.gov/oppde/larc">www.fsis.usda.gov/oppde/larc</a> at the "ingredients" link)

## 5. Question: What is the maximum amount of organic acid(s) permitted to be applied to livestock carcasses pre-chill without having to declare the organic acid(s) on the label?

Answer: Historically, the maximum amount of organic acid(s) that can be used to treat livestock carcasses without labeling is up to 2.5 % of a solution applied pre-chill. Labeling is not required for this specific use of organic acid(s) (which the Agency has permitted for many years) because it is based on data that showed that this application is consistent with FDA's definition of a processing aid.

Recently, FSIS has recently stated no objection to the use of 5 % hot lactic acid as an antimicrobial agent on beef carcasses prior to fabrication (see question number four). This use was determined to be consistent with the definition of a processing aid. Therefore, its use would not need to be reflected on the labeling for treated carcasses or products produced from treated carcasses.

## 6. Question: Do organic acid(s) (e.g., lactic, acetic, or citric acid) that are used as antimicrobial agents need to be declared on the label if they are applied to livestock carcasses?

Answer: Unless the proposed use has been determined by FSIS to be consistent the definition of a processing aid (e.g., the application of acetic or citric acids at 2.5 % of a beef carcass wash solution applied pre-chill or the use of a 5% lactic acid solution to treat beef carcasses prior to fabrication either pre- or post-chill) the organic acid(s) would require labeling.

# 7. Question: Is the maximum amount of organic acid(s) allowed, without labeling the product, based on the concentration of the organic acid(s) applied to the carcass or the concentration of the organic acid(s) draining from the carcass?

Answer: The amount of organic acid(s) is based on the percentage of organic acid(s) in the carcass wash (aqueous solution) prior to application. It is not based on the residual level of organic acid(s) draining from a treated carcass during application.

### 8. Question: Do organic acid(s) (e.g. lactic, acetic, or citric acid) have to be declared on the label if they are applied to cut-up and ground meat and poultry?

Answer: Yes, all ingredients, including organic acid(s), require labeling unless the use of a substance is consistent with FDA's definition of a processing aid or is a secondary direct food additive. If an establishment is interested in using organic acid(s) to treat meat and poultry cuts and/or ground meat and poultry to momentarily reduce microorganisms, data must be submitted to FSIS to show that the proposed use of organic acid(s) is consistent with FDA's definition of a processing aid.

# 9. Question: Do organic acid(s) (e.g. lactic, acetic, or citric acid) have to be declared on the label if they are applied to livestock or poultry byproducts and giblets (e.g. livers, hearts, and gizzards)?

Answer: No, labeling is not required when organic acid(s) are applied pre-chill at up to 2.5% of an aqueous solution to treat livestock and poultry byproducts and giblets.

FSIS has recently stated no objection to the use of 5% lactic acid as an antimicrobial agent to treat beef carcasses prior to fabrication (i.e., pre and post-chill).

## 10. Question: Are organic acid(s) used as antimicrobial agents permitted to be used on poultry carcasses?

Answer: Yes, organic acid(s) are GRAS and are listed in FSIS regulations for use as an acidifier (which may have an antimicrobial effect) in various meat and poultry products at a level which is sufficient for purpose (9 CFR 424.21(c)). Organic acid(s) are permitted to be applied to poultry carcasses pre-chill at a concentration of up to 2.5 percent of a solution without labeling.

### 11. Question: If organic acid(s) (e.g., lactic, acetic, or citric acid) are used on ready-to-eat products as a spray or dip, must the application be followed by a potable water rinse?

Answer: No, the use of organic acid(s) on ready-to-eat products are not required to be followed by a potable water rinse. However, the organic acid(s) will be considered ingredients that require labeling unless data can be submitted to FSIS that show that their use is consistent with FDA's definition of a processing aid.

12. Question: Are organic acid(s) (e.g., lactic, acetic or citric acid) permitted to be used on a continuous basis on conveyor belts? What are the conditions for their use? When do the organic acids need to be declared on a product label?

Answer: FSIS has no objection to the use of organic acids on conveyor belts on a continuous basis. However, the process should not result in the organic acid(s) having a lasting technical effect on meat or poultry which comes into contact with the conveyor belts. Labeling is required if the organic acid(s) exhibit a lasting technical effect on meat or poultry which comes into contact with the treated conveyor belts.

13. Question: Are antimicrobial agents other than organic acid(s) permitted to be used on a continuous basis on conveyor belts if they are approved as an antimicrobial agent in the production of meat and poultry products? What are the conditions for their use? When do the antimicrobial agents have to be included on a product label?

Answer: Yes, antimicrobial agents approved for use in the production of meat and poultry products may be used on conveyor belts provided they are followed by a potable water rinse. Substances listed in 21 CFR 178.1010 may be used in sanitizing solutions on food contact surfaces with only adequate draining (no water rinse) before contact with food.

14. Question: Is trisodium phosphate (TSP) permitted to be used as an antimicrobial agent on livestock carcasses, viscera, and parts?

Answer: TSP may only be used on livestock carcasses according to interim Agency policy.

15. Question: Where is TSP allowed to be used as an antimicrobial agent on poultry?

Answer: FSIS regulations (9 CFR 424.21 (c)) permits the use of TSP on raw post-chill poultry carcasses. In addition, FSIS has permitted the application of TSP to raw poultry carcasses pre-chill by spraying or dipping the carcasses with an 8-12% solution maintained within a temperature range of 65° F to 85° F for up to 15 seconds. FSIS has permitted the use spraying or dipping of poultry giblets for up to 30 seconds with an 8-12% solution of TSP pre-chill.

TSP is also used in some on-line reprocessing operations. Establishments which use on-line reprocessing operate under an experimental exemption listed in 9 CFR 381.3(c). The conditions of use for TSP in on-line reprocessing are limited by the parameters listed in the FSIS approved on-line reprocessing protocol, not the conditions of use listed above.

16. Question: Is chlorine dioxide permitted to be used as an antimicrobial agent on livestock carcasses, viscera, and parts?

Answer: Chlorine dioxide may be used as an antimicrobial agent to treat red meat carcasses, parts, and organs. It is applied as a spray or dip at a level not to exceed 3 ppm residual chlorine dioxide.

17. Question: Is chlorine dioxide allowed to be used as an antimicrobial agent on poultry? What are the conditions for its use?

Answer: Chlorine dioxide may be used as an antimicrobial agent to treat water in poultry processing as prescribed in FDA's regulations (21 CFR 173.300). Residual chlorine dioxide must not exceed 3 ppm in the poultry processing water.

### 18. Question: Is hydrogen peroxide allowed to be used as an antimicrobial agent on meat and poultry products (e.g. carcasses, parts, processed products)?

Answer: No, hydrogen peroxide is listed as GRAS in FDA regulations (21 CFR 184.1366) for use as a bleaching agent to treat beef feet and in FSIS regulations (9 CFR 424.21 (c)) as a bleaching agent to treat tripe (followed by a water rinse). It is also a component of peroxyacids (21 CFR 173.370).

### 19. Question: Can any and all antimicrobial agents be used on poultry carcasses during online reprocessing?

Answer: No, on-line reprocessing operations function under an experimental exemption (9 CFR 381.3 (c)). The use of antimicrobial agents in on-line reprocessing are limited by the parameters of the FSIS approved on-line reprocessing protocol.

## 20. Question: Can antimicrobial agents be used (spray or dip) on the same carcasses or parts more than once, without labeling?

Answer: Yes, antimicrobial agents may be used more than once. However, the antimicrobial agents must be used in accordance with the approved or accepted conditions of use. Labeling is required unless the use of the substance is consistent with FDA's definition of a processing aid or is a secondary direct food additive.

## 21. Question: Do all uses of antimicrobial agents need to comply with the requirements of 9 CFR 441.10 for retained water? What are the requirements?

Answer: Yes, any establishment that uses a post-evisceration process that results in water retention in raw livestock or poultry carcasses or parts must maintain on file a written data collection protocol in accordance with 9 CFR 441.10 (c) (1). Any treatment in the chilling process such as antimicrobial treatments should be described in the protocol. An establishment does not have to maintain a protocol on file if it has data or information that clearly demonstrates that its products do not retain water as a result of the process, e.g., spraying boneless meat with antimicrobial agents where the end product does not retain water from the antimicrobial application <u>FSIS Directive 6700.1</u> and <u>6700.1 Amend 1</u>).